

Is Uterine Compression Suture for Postpartum Hemorrhage safe? A Case report

Abstract

The main cause of PPH is uterine atony. Uterine compression suturing is a common technique to control post-partum hemorrhage (PPH) at cesarean delivery. Here, we report a complication of this method for atony post-delivery.

A primigravida 27 years woman with term pregnancy underwent cesarean delivery and because of uterine atony unresponsive to medical therapy and uterine artery ligation so two compression sutures were placed on the uterus. But after 11 days, our patient underwent surgery Again, due to severe fever, infection and necrotic mass in the uterine cavity. During operation we removed the necrotic mass.

Although Uterine Compression Suture is an effective method for the treatment of PPH, in our study we have reported some side effects specially necrosis of myometrium in this technique.

Keywords: Postpartum hemorrhage, surgical treatment, Suture Techniques, uterine atony

Introduction

Postpartum hemorrhage (PPH) is one of the most common causes of maternal morbidity and mortality all over the world(1). The main cause of PPH is uterine atony. The treatment of uterine atony includes medical therapy and surgical treatment (2). Among different techniques to manage the PPH, Uterine compression sutures -especially B-lynch suture- have most popularity and many studies have evaluated its efficacy. Although most studies have shown no complications for compression sutures, in this report, we present a case of uterus

atony treated with b-lynch suture which has complicated with postoperative abscess inside the myometrium.

Case Report

A primigravida 27 years woman with term pregnancy in active phase of labor was admitted. On admission to hospital, cervical examination included dilatation 5cm, effacement 60% and station -2. Vital signs were normal and she had no special medical and surgical history. Dilatation became full within 3 hours with spontaneous contractions, but after 2 hours she underwent cesarean section due to transverse arrest. Because of mild atony during cesarean we administered high dose of oxytocin(100 unit in a 1liter ringer lactate) and rectal misoprostol(800 microgram) .We closed the abdominal wall and patient was transferred to the recovery with good general status. After 3 hours of complete care she was sent to the obstetrics ward.

The patient had again uterine atony and vaginal bleeding 6 hours after the operation .Despite uterus massage and readministration of oxytocin and misoprostol, haemorrhage continued so she was candidate for relaparotomy 9 hours after cesarean. Due to no response and continuing bleeding, uterine was opened by kerr incision. The uterine was packed with a long gauze and again b-lynch suture and a cyclic suture in uterine isthmus were placed to preserve the uterine.

After bleeding cessation, the abdominal wall was closed and the patient was admitted to the Intensive Care Unit (ICU) with order of cefazolin and gentamycin. In the ICU, the patient's general condition was good and she had no vaginal bleeding. The patient's tests in the ICU included: platelet=30000, white blood cell=21.000, Alanine transaminase (ALT) =260, aspartate transaminase(AST)=300, INR=4, creatinine=1.Long gauze was removed through the vagina 36 hours after the surgery.

After 3days, regarding to high fever, antibiotics including meropenem and vancomycin were

started for her. Breast and abdominal examinations and chest X-ray were normal. In blood culture no organism was grown.

All patients' tests were improved 6 days after surgery and the patient was discharged without fever and antibiotic. 5 days after discharge the patient was rehospitalized with a high fever (T=40°C), PR=130, BP=100/60. Fundal height was 24 weeks. In bimanual examination she had irregular and tender uterus. Cervix was open, but did not have vaginal bad smell discharge. X-Ray of the abdomen and pelvis were performed to confirm the absence of external object.

Computed Tomography(CT) scan of the abdomen and pelvis reported large uterus and Pear Shape structure with liquid air density that continues to the pelvic floor. In transvaginal ultrasound, we observed some gases in the myometrium and the liquid air level inside the uterus so the patient underwent curettage by guided-sonography. Although the uterus cavity was empty, Brief necrotic myometrial tissue was removed.

According to diagnosis of abscess inside the myometrium, intravenous antibiotics including ampicillin-sulbactam, meropenem, and metronidazole were started for her. After 2 days, fever was stopped and she was discharged with a recommendation to continue treatment for up to 3 weeks. Two weeks after discharge, the patient referred to the clinic due to T = 38°C.

In vaginal examination, there was necrotic mass with the approximate dimensions of 10x 12cm that was exited from the cervix and was torn during the examination and about 400-300cc smelly pus came out. The results of gram-negative and gram-positive cocci were reported. The patient did not consent to hospitalization and underwent antibiotic experimental regimen (metronidazole). Finally, the patient was examined 4 weeks later when the uterine was normal and there was also no abnormality in the sonography of uterine and ovaries.

Discussion

Post-Partum Hemorrhage (PPH) causes approximately 19.1% of mother's death after delivery, for example in France has reported 1.6 death/100,000 live births(7).The prevalence of PPH has increased 27.5% from 1995 to 2004. Primarily the main reason of this increase is high incidence of uterine atony(8).So surgical treatment like compression sutures is used after the failure of medical therapy. Studies have reported the success rate of 91.7% for a variety of compression sutures(9) while according our recent study, We believe that, advantages and disadvantages of therapeutic options even compression sutures should be monitored.

Probably achievement to successes in compression sutures is related to the protective physiological process in uterus. This means that it can be explanted in order to rapid involution of uterus within the first postpartum week(10).

Correspondingly Tsitlakidis et al have presented a case with successful pregnancy after PPH during longest follow up (10 years), other Published data have certified that the B-Lynch surgical technique is secure, advantageous and with lack of short and long-term complication.(11) However, there are concerns about this technique that include uterine adhesions, partial ischemia, necrosis, infection, and myometra(9).

For example, Akoury Het alhas reported A healthy 32-year-old woman, gravida 3, para 0. In order to control PPH, They were used 2 Cho and 1 B-Lynch sutures. Subsequently they have identified a large triangular myometrial defect and two smaller defects respectively in the mid-anterior and in the posterior uterine wall (12). Also in the study of Gottlieb AGa 33-year gravida 2, para 0101 due to uterine atony, received two compression sutures. On the 8th day of post-operative because of high fever antibiotic-resistant, patient was sent to the operating room for a second time. During a laparotomy they found fundal uterine necrosis in suture place(13). Reyftmann L has noted a partial uterine necrosis after placement Of Four Cho

sutures for control of PPH(14). However, our patient underwent surgery Again, due to severe fever, infection and necrotic mass in the uterine cavity. During operation we removed the necrotic mass.

Various materials are used for sutures like vicryl, PDS and nylon. Studies have emphasized on avoiding the usage of none or slowly absorbable sutures(15). Wu HH et al in their study have placed multiple square sutures for a primigravida with severe PPH .Consequently they realized partial obstruction of menstrual flow with uterine synechiae. Hysteroscopy revealed unabsorbed sutures in uterine (16). In our study, for a primigravida woman we have used sutures. So soluble sutures should be used to reduce the risk of uterine rupture and necrosis in this procedure, however uterine rupture has been reported during manual compression sutures(17).

Sentilhes et all have observed a case of Asherman syndrome with obliteration of the uterine cavity after placement of B- Lynch sutures, which may lead to temporary or permanent infertility(18). Several cases of successful pregnancy after compression sutures techniques have been introduced(19). It is necessary that physicians and patients be aware of the possible complications of using this technique(12).

So we suggest the postoperative follow-up to establish a national registry of cases with placement of compression sutures. Then we can document the effect of this method on potential fertility, infertility, menstrual flow, and evaluated the long and short-term complications of this procedure.

However, it is difficult to observe the fertility outcomes of women with compression sutures, because some of them avoid pregnancy due to fear of recurring PPH(19, 20). There is also limited data on women's menstrual period. These defects reveal the need of a database to register the success or failure results. Moreover possible complications of compression suture such as B-Lynch suture will be recorded (9).

In summary, most studies have pointed to high efficiency and absence of side effects in compression suture such as B-Lynch suture. While in our study and some other similar studies have reported more side effects such as high fever resistant to antibiotics, ischemia and necrosis of the uterus, uterine adhesions, and etc.

According to inconsistencies, it is necessary to follow the women who were underwent compression sutures. Besides prospective studies is recommended. Therefore we can collect reliable database, in order to assess the short and long term effects of this technique, and then to answer critical questions that: Is it affects the future potential fertility?

Conclusion

Although Uterine Compression Suture is an effective method for the treatment of PPH, The surgeon should be aware of side effects of this technique.

References

1. Pereira A, Nunes F, Pedroso S, Saraiva J, Retto H, Meirinho M. Compressive uterine sutures to treat postpartum bleeding secondary to uterine atony. *Obstet Gynecol.* 2005;**106**:569-72.
2. Ochoa M, Allaire AD, Stitely ML. Pyometria after hemostatic square suture technique. *Obstet Gynecol.* 2002;**99**:506-9.
3. Baskett TF. Uterine compression sutures for postpartum hemorrhage: efficacy, morbidity, and subsequent pregnancy. *Obstet Gynecol.* 2007;**110**:68-71.
4. Halder A. A new uterine suture technique to control PPH in congenitally malformed uterus during caesarean section. *Obstet Gynecol.* 2009;**29**:402-4.
5. Coker A, Lawal AH, Abu J, Cowen MJ. The B-Lynch surgical technique for the control of massive postpartum haemorrhage: an alternative to hysterectomy? Five cases reported. *BJOG.* 1997;**104**:372-5.

6. Zheng J, Xiong X, Ma Q, Zhang X, Li M. A new uterine compression suture for postpartum haemorrhage with atony. *BJOG*. 2011;**118**:370-4.
7. Deneux-Tharoux C, Bonnet M, Tort J. [Epidemiology of post-partum haemorrhage.]. *J Gynecol Obstet Biol Reprod*. 2014;**43**:936-50.
8. Bateman BT, Berman MF, Riley LE, Leffert LR. The epidemiology of postpartum hemorrhage in a large, nationwide sample of deliveries. *Anesth Analg*. 2010;**110**:1368-73.
9. Saroja CSM, Nankani A, El-Hamamy E. Uterine compression sutures, an update: review of efficacy, safety and complications of B-Lynch suture and other uterine compression techniques for postpartum haemorrhage. *Arch Gynecol Obstet*. 2010;**281**:581-8.
10. Joshi VM, Shrivastava M. Partial ischemic necrosis of the uterus following a uterine brace compression suture. *BJOG*. 2004;**111**:279-80.
11. Tsitlakidis C, Alalade A, Danso D, C BL. Ten year follow-up of the effect of the B-Lynch uterine compression suture for massive postpartum hemorrhage. *Int J Fertil Womens Med*. 2006;**51**:262-5.
12. Akoury H, Sherman C. Uterine wall partial thickness necrosis following combined B-Lynch and Cho square sutures for the treatment of primary postpartum hemorrhage. *JOGC* . 2008;**30**:421-4.
13. Gottlieb AG, Pandipati S, Davis KM, Gibbs RS. Uterine necrosis: a complication of uterine compression sutures. *Obstet Gynecol*. 2008; **112**:429-31.
14. Reyftmann L, Nguyen A, Ristic V, Rouleau C, Mazet N, Dechaud H. [Partial uterine wall necrosis following Cho hemostatic sutures for the treatment of postpartum hemorrhage]. *Gynecol Obstet Fertil*. 2009;**37**:579-82.
15. Cotzias C, Girling J. Uterine compression suture without hysterotomy--why a non-absorbable suture should be avoided. *J Obstet Gynaecol*. 2005;**25**:150-2.

16. Wu HH, Yeh GP. Uterine cavity synechiae after hemostatic square suturing technique. *Obstet Gynecol.* 2005;**105**:1176-8.
17. Higgins L, Chan K, Tower C. Uterine rupture following previous uterine compression suture. *J Obstet Gynaecol.* 2011;**31**:544.
18. Sentilhes L, Trichot C, Resch B, Sergent F, Roman H, Marpeau L, et al. Fertility and pregnancy outcomes following uterine devascularization for severe postpartum haemorrhage. *Hum Reprod.* 2008;**23**:1087-92.
19. Goojha CA, Case A, Pierson R. Development of Asherman syndrome after conservative surgical management of intractable postpartum hemorrhage. *Fertil Steril.* 2010;**94**:1098 e1-5.
20. Treloar E, Anderson R, Andrews H, Bailey J. Uterine necrosis following B-Lynch suture for primary postpartum haemorrhage. *BJOG.* 2006;**113**:486-8.