

Investigation of the Influence of Serum Selenium Levels on the Risk of Laryngeal Cancer

Simin Akbari¹, Bashir Rasoulia¹ , Imaneh Roshan Zamir¹ , Mona Kabiri² , Kiana Ketabi³ ,
 Ehsan Khadivi^{1*} 

¹ Sinus and Surgical Endoscopic Research Center, Ghaem Hospital, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

² Nanotechnology Research Center, Pharmaceutical Technology Institute, Mashhad University of Medical Sciences, Mashhad, Iran

³ Department of Otorhinolaryngology, Sinus and Surgical Endoscopic Research Center, School of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

⁴ Sinus and Surgical Endoscopic Research Center, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

ARTICLE INFO

Article type

Original article

Article history

Received: 05 March 2025

Accepted: 16 Nov 2025

Keywords

Laryngeal cancer

Selenium

Aerodigestive system

Case-control study

ABSTRACT

Introduction: Cancers of the upper aerodigestive tract are relatively common, with laryngeal malignancies ranking second in prevalence. Selenium is an essential trace element whose biological functions in human cells remain incompletely elucidated. While moderate selenium levels may provide protective antioxidant effects, both deficiency and excess have been associated with carcinogenic processes. The present study aimed to evaluate the relationship between serum selenium concentration and the likelihood of laryngeal cancer development.

Methods: We conducted a case-control study involving 30 untreated patients with laryngeal cancer and 30 disease-free individuals, recruited from Ghaem and Imam Reza Hospitals in Mashhad, Iran. The two groups were matched for age, sex, cigarette exposure (pack-years), duration of opium consumption, symptoms of gastroesophageal reflux, and occupational dust exposure. Venous blood samples were collected before treatment to measure serum selenium levels, and mean values were statistically compared between cases and controls.

Results: Across all 60 participants, the mean serum selenium level was 96.37 ± 20.13 $\mu\text{g/L}$. Patients in the cancer group exhibited higher concentrations (102.89 ± 19.40 $\mu\text{g/L}$) than those in the control group (89.85 ± 18.98 $\mu\text{g/L}$). This difference was statistically significant ($p = 0.011$).

Conclusion: This study indicated that elevated serum selenium levels may be linked with an increased probability of laryngeal cancer.

Please cite this paper as:

Akbari S, Rasoulia B, Roshan Zamir I, Kabiri M, Khadivi E. Investigation of the Influence of Serum Selenium Levels on the Risk of Laryngeal Cancer. *Reviews in Clinical Medicine*. 2025;12(4): 126-130.

Introduction

Laryngeal cancer is the second most common malignancy of the upper aerodigestive tract, with squamous cell carcinoma (SCC) representing the predominant histological type, accounting for approximately 85%–95% of all laryngeal tumors. A

large-scale review of nearly 160,000 laryngeal SCC cases in the United States showed that the glottis was the primary site in 51% of patients, the supraglottis in 33%, and the subglottis in 2%, while in 14% of cases the exact location could not be determined. The

***Corresponding author:** Dr. Ehsan Khadivi, Sinus and Surgical Endoscopic Research Center, Ghaem Hospital, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran
Email: khadivie@mums.ac.ir

Doi: [10.22038/RCM.2025.87973.1544](https://doi.org/10.22038/RCM.2025.87973.1544)

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

overall male-to-female incidence ratio for laryngeal carcinoma is estimated at 3.9:1 (1, 2).

Tobacco use and alcohol consumption remain the most significant risk factors for the development of laryngeal malignancies (3), with risk increasing in proportion to both the duration and intensity of exposure. Chronic laryngeal irritation has also been proposed as a potential etiological factor, particularly among individuals without a history of smoking or alcohol use (4). Occupational exposure to hazardous substances constitutes another critical contributor. Incidence rates are higher among manual laborers who not only consume greater amounts of tobacco and alcohol but are also frequently exposed to toxic agents. Reported occupational hazards include diesel fumes, asbestos, organic solvents, sulfuric acid, mustard gas, mineral oils, metal dusts, asphalt, wood and stone dust, mineral wool, and cement particles. However, establishing a causal relationship between any specific toxin and laryngeal cancer has been challenging due to small sample sizes and multiple confounding variables in most studies (5). Human papillomavirus (HPV) is a well-established causative factor in cervical cancer and is increasingly recognized as a key etiological agent in oropharyngeal squamous cell carcinoma (SCC) (6). Moreover, genetic predisposition to laryngeal SCC should not be overlooked (7). Evidence from a cohort study reported a relative risk of 3.79 for head and neck SCC among individuals with a positive family history of these malignancies (2). Dietary patterns also influence risk; adherence to a diet rich in fruits and vegetables and lower in meat and fat has been associated with a decreased incidence of head and neck cancers (8).

Selenium is an essential trace element involved in multiple biological processes, including antioxidant defense and immune regulation, and is incorporated into at least 30 selenoproteins (9). These proteins contribute to maintaining redox homeostasis and preserving DNA integrity, and some studies suggest that they may exhibit anti-carcinogenic properties (10). Selenium also helps neutralize the toxic effects of heavy metals and other harmful agents. Clinical evidence has indicated that selenium supplementation may reduce the risk of certain cancers. Notably, both selenium deficiency and excessive intake have been associated with an increased incidence of malignancies such as lung, laryngeal, colorectal, and prostate cancers (11, 12).

This case-control study aims to compare serum selenium levels between patients with laryngeal

cancer and a matched control group to assess whether selenium status is associated with the risk of developing laryngeal cancer.

Materials

Patients and Methods

This case-control study was conducted from December 2022 to April 2023 on 30 newly diagnosed patients with laryngeal cancer from the ENT departments of Imam Reza and Ghaem Hospitals, along with a matched group of 30 healthy controls. The two groups were comparable based on age, sex, and smoking status. Participation required written informed consent; individuals who declined were not enrolled.

The case group included patients with a recent histopathological diagnosis of laryngeal cancer who were enrolled before the initiation of any treatment. The control group consisted of individuals with no history of malignancy and no use of selenium supplements within the month before enrollment. Given the high prevalence of smoking and opium use among patients with laryngeal cancer, controls were selected through convenient non-random sampling from patients admitted to the cardiopulmonary wards of Imam Reza and Ghaem Hospitals who reported a history of smoking.

Exclusion criteria included withdrawal from participation, a prior diagnosis of malignancy, recurrent disease, or selenium supplementation during the study period. For all patients, the diagnosis was confirmed histopathologically, and tumor mapping was performed to determine the histological subtype. Baseline demographic and clinical information was collected using a structured questionnaire and included variables such as age, sex, tobacco and alcohol use, symptoms of gastric reflux, family history of cervical cancer, degree of familial relation, occupation, and exposure to stone or wood dust.

Following data collection, 3 mL of venous blood was drawn from each participant. Serum specimens were then processed and analyzed using a graphite furnace atomic absorption spectrophotometer with selenium-specific wavelengths. Calibration was performed using standard selenium solutions, and quality control samples were run alongside each batch to ensure accuracy and precision. The measured absorbance values were converted to selenium concentrations according to the calibration curve. Selenium levels were subsequently compared between cases and controls. All data were analyzed using SPSS software.

Statistical Analysis

Data were analyzed using SPSS version 20. Sample size was estimated with PASS version 15 using a two-sample t-test assuming unequal variances. A minimum of 28 participants per group was required with a significance level (α) of 0.05 and a power ($1-\beta$) of 90%; therefore, 30 individuals were included in each group.

Continuous variables were summarized as mean \pm standard deviation. The normality of the data distribution was assessed using the Kolmogorov-Smirnov test. Depending on the distribution, group comparisons for quantitative variables were performed using the independent t-test. Categorical variables were presented as frequencies and percentages and analyzed using the chi-square test or Fisher's exact test, as appropriate. A p-value of less than 0.05 was considered statistically significant.

Results

In this case-control study, 60 participants were enrolled. The case group consisted of 30 patients diagnosed with laryngeal squamous cell carcinoma who were evaluated before receiving any treatment (chemoradiotherapy or surgery). The control group included 30 individuals without

laryngeal cancer who were matched to the cases based on gender, age, and the extent of smoking and opium consumption.

The mean age of participants in the control group was 59.86 ± 11.67 years, whereas in the case group it was 57.80 ± 9.93 years. There was no significant difference in age between the two groups ($p \geq 0.05$). According to the Kolmogorov-Smirnov test, the data for serum selenium levels and age followed a normal distribution.

Among the 60 participants, each group consisted of 28 men (93.3%) and two women (6.7%). In both groups, 27 individuals were smokers and 3 were non-smokers. Regarding opium consumption, 26 patients in the case group (86.7%) and 23 patients in the control group (76.7%) reported opium use, with no significant difference between the groups ($p = 0.506$; $p \geq 0.05$). None of the participants had a history of alcohol consumption.

All patients were asked about symptoms of esophageal reflux, and no significant difference was observed between the two groups ($p = 0.299$; $p \geq 0.05$). Regarding occupation, 73.3% of participants in both groups were manual laborers (including construction and agricultural workers), with no significant difference between the groups ($p > 0.9$) (Table 1).

Table 1. Number and percentage of subjects with a history of contact with dust and GERD (Gastro Esophageal Reflux Disease) in the case and control groups

			Control group	Laryngeal cancer group	P-value*
Contact with dust	No	Number	8	8	>0.9
		Percent	26.7%	26.7%	
Yes	Number	22	22		
	Percent	73.3%	73.3%		
GERD	No	Number	19	14	0.299
		Percentage	63.3%	46.7%	
	Yes	Number	11	16	
		Percentage	36.7%	53.3%	
Total		Number	30	30	

Chi-Square test*

We classified all 30 patients in the laryngeal cancer group by tumor stage into early-stage (T1, T2) and advanced-stage (T3, T4) disease. In addition, the tumor anatomical subsite was categorized into three subtypes: supraglottic, glottic, and transglottic tumors (Figure 1). Analyses using the Pearson correlation test showed no significant relationship between tumor characteristics and serum selenium levels.

The mean serum selenium level among all 60 participants was 96.37 ± 20.13 $\mu\text{g/L}$, while the normal laboratory reference range is 87–154 $\mu\text{g/L}$.

A comparison of mean serum selenium levels between the laryngeal cancer group (before treatment) and the control group is presented in Figure 2. The mean serum selenium level was 102.89 ± 19.40 $\mu\text{g/L}$ in the laryngeal cancer group and 89.85 ± 18.98 $\mu\text{g/L}$ in the control group. The findings indicated that serum selenium levels were significantly higher in the laryngeal cancer group compared with the control group ($p = 0.011$; $p < 0.05$).

Discussion

The findings of this case-control study, which compared 30 patients with laryngeal cancer with 30 matched hospital-based controls, showed no significant association between age or tumor stage and serum selenium levels. Notably, higher serum selenium levels were associated with a higher risk of laryngeal cancer.

To our knowledge, only a limited number of studies have investigated selenium status in relation to laryngeal cancer progression or clinical outcomes, and the available evidence remains inconsistent. Such discrepancies may reflect the dual biological roles of selenium in normal and malignant cells. For example, sodium selenite has been shown to promote apoptosis in lung cancer cell lines (13), whereas other studies have suggested a potential protective effect against the development of aggressive tumors (14).

Consistent with our findings, Narod et al conducted a case-control study in Poland titled "Evaluation of serum selenium levels and cancer risk." The study included 97 women with various types of cancer and 184 cancer-free controls at baseline. Using mass spectroscopy, the authors defined normal serum selenium levels as 70–90 µg/L. Selenium levels above 90 µg/L were associated with an increased overall cancer risk, although this association did not reach statistical significance. Their findings indicated a non-linear relationship between selenium status and cancer susceptibility (15-16). Similarly, Krystal et al evaluated selenium exposure using nail samples from more than 3,000 American men to examine prostate cancer risk. Their findings indicated that selenium supplementation did not confer protection among men with low baseline selenium levels; however, higher selenium status was associated with an increased likelihood of high-grade prostate cancer in men with elevated selenium levels (16).

Comparable findings were also reported in Iran by Shahriari et al, who investigated soil selenium levels in Golestan Province. Selenium concentrations were measured across 135 soil blocks, yielding an average value of 1.61 ± 3.7 mg/kg. Higher selenium levels were observed in high-risk regions for esophageal cancer (4.13 mg/kg) compared with lower-risk areas (3.39 mg/kg). The authors concluded that soil selenium levels may directly contribute to the regional burden of esophageal cancer (17).

Conversely, Jaworska et al reported differing findings in their Polish case-control study titled "Low selenium level is associated with laryngeal

cancer." The study included 95 patients with lung cancer, 113 patients with laryngeal cancer, and matched healthy controls. Selenium levels were lower in both cancer groups compared with their respective control groups. Specifically, the mean selenium concentration was 63.2 µg/L in lung cancer patients versus 74.6 µg/L in controls, and 64.8 µg/L in laryngeal cancer patients versus 77.1 µg/L in their matched controls. The authors proposed that selenium concentrations below 60 µg/L may increase the risk of both lung and laryngeal cancer (18).

In contrast, our results showed higher mean selenium levels in both groups: 102.89 ± 19.40 µg/L in patients and 89.85 ± 18.98 µg/L in controls. These values exceed those reported by Jaworska et al., suggesting that participants in our study were not, on average, selenium-deficient. This discrepancy may partly account for the differences observed between our findings and theirs.

Furthermore, selenium status varies geographically. For instance, the mean selenium intake is generally higher in North America than in many European countries. Large randomized controlled trials conducted in populations with adequate baseline selenium intake, such as those in the United States and Canada, have not demonstrated a clear reduction in cancer incidence or mortality with selenium supplementation. In contrast, studies from regions with relatively low selenium status, including Poland and Sweden, have suggested that higher selenium levels may improve survival in certain cancers, such as breast cancer.

Conclusions

We concluded that studies reporting an inverse relationship between serum selenium levels and cancer risk were predominantly conducted in countries with relatively low normal selenium levels. In contrast, the present study and similar investigations conducted in Iran indicate that our population has comparatively high average selenium levels. Therefore, given this higher baseline status, the hypothesis that selenium deficiency contributes to the development of laryngeal cancer was not supported by our findings.

Ethics approval and consent to participate

The present study was approved by Research Council of Mashhad University of Medical Sciences and the ethics committee approval code number is (IR.MUMS.MEDICAL.REC.1401.561). The written informed consent forms were signed by all patients before initiation of the study.

Consent for publication

Complete written informed consent was obtained from the patient for the publication of this study

Availability of data and materials

The dataset supporting this article's conclusions is confidential and cannot be made publicly available. However, data may be available upon reasonable request and with permission of the original data providers, subject to compliance with applicable confidentiality agreements and data protection laws, via contact through the corresponding author.

Competing interests

The authors declare that there is no conflict of interest regarding the publication of this paper.

Funding

This research was financially supported by vice president for research, Mashhad University of Medical Sciences, Mashhad, Iran (grant number: 4000574). These data have been extracted from the thesis of simin Akbari, to obtain Specialized assistant degree, Medical School, Mashhad University of Medical Sciences.

Authors' contributions

Writing original draft: Simin Akbari, Kiana Ketabi, Ehsan Khadivi

Reviewing: Bashir Rasoulia, Imaneh Roshan Zamir, Mona Kabiri

Conceptualization: Ehsan Khadivi, Simin Akbari, Kiana Ketabi

Acknowledgements

Thanks to the deputy or research, Mashhad University of Medical Sciences.

References

- Lubiński J, Marciniak W, Muszynska M, Jaworowska E, Sulikowski M, Jakubowska A, et al. Serum selenium levels and the risk of progression of laryngeal cancer. *PLoS One*. 2018;13(1):e0184873. doi:10.1371/journal.pone.0184873 PMID:29304040
- Kraft S, Schindler J. Cummings otolaryngology head and neck surgery: Tracheotomy. Flint PW, editor. Philadelphia: Elsevier Saunders; 2020.
- Kulháňová I, Forman D, Vignat J, Espina C, Brenner H, Storm HH, et al. Tobacco-related cancers in Europe: The scale of the epidemic in 2018. *European Journal of Cancer*. 2020;139:27-36. doi:10.1016/j.ejca.2020.07.024 PMID:32957011
- Kost KM, Jefferson GD. Larynx Cancer. *Otolaryngologic Clinics of North America*. 2023;56(2):xix-xx. doi:10.1016/j.otc.2023.02.001 PMID:37030952
- Hashibe M, Brennan P, Benhamou S, Castellsague X, Chen C, Curado MP, et al. Alcohol drinking in never users of tobacco, cigarette smoking in never drinkers, and the risk of head and neck cancer: pooled analysis in the International Head and Neck Cancer Epidemiology Consortium. *Journal of the National Cancer Institute*. 2007;99(10):777-89.

- doi:10.1093/jnci/djk179 PMID:17505073
- Yang D, Shi Y, Tang Y, Yin H, Guo Y, Wen S, et al. Effect of HPV infection on the occurrence and development of laryngeal cancer: a review. *Journal of Cancer*. 2019;10(19):4455. doi:10.7150/jca.34016 PMID:31528209
 - de Miguel-Luken MJ, Chaves-Conde M, Carnero A. A genetic view of laryngeal cancer heterogeneity. *Cell Cycle*. 2016;15(9):1202-12. doi:10.1080/15384101.2016.1156275 PMID:26940775
 - Sittitrai P, Ruenmarkkaew D, Booyaprapa S, Kasempitakpong B. Effect of a perioperative immune-enhancing diet in clean-contaminated head and neck cancer surgery: a randomized controlled trial. *International Journal of Surgery*. 2021;93:106051. doi:10.1016/j.ijssu.2021.106051 PMID:34371175
 - Minich WB. Selenium metabolism and biosynthesis of selenoproteins in the human body. *Biochemistry (Moscow)*. 2022;87(Suppl1):S168-S77. doi:10.1134/S0006297922140139 PMID:35501994
 - Szwiec M, Marciniak W, Derkacz R, Huzarski T, Gronwald J, Cybulski C, et al. Serum selenium level predicts 10-year survival after breast cancer. *Nutrients*. 2021;13(3):953. doi:10.3390/nu13030953 PMID:33809461 PMID:PMC7998294
 - Pietrzak S, Wójcik J, Scott RJ, Kashyap A, Grodzki T, Baszuk P, et al. Influence of the selenium level on overall survival in lung cancer. *Journal of Trace Elements in Medicine and Biology*. 2019;56:46-51. doi:10.1016/j.jtemb.2019.07.010 PMID:31442953
 - Al-Ansari RF, Al-Gebori AM, Sulaiman GM. Serum levels of zinc, copper, selenium, and glutathione peroxidase in the different groups of colorectal cancer patients. *Caspian Journal of Internal Medicine*. 2020;11(4):384. doi:10.22088/cjim.11.4.384
 - Jönsson-Videsäter K, Björkhem-Bergman L, Hossain A, Söderberg A, Eriksson LC, Paul C, et al. Selenite-induced apoptosis in doxorubicin-resistant cells and effects on the thioredoxin system. *Biochemical pharmacology*. 2004;67(3):513-22. doi:10.1016/j.bcp.2003.09.021 PMID:15037203
 - Heras IL, Palomo M, Madrid Y. Selenoproteins: the key factor in selenium essentiality. State-of-the-art analytical techniques for selenoprotein studies. *Analytical and bioanalytical chemistry*. 2011;400:1717-27. doi:10.1007/s00216-011-4916-4 PMID:21455649
 - Narod SA, Huzarski T, Jakubowska A, Gronwald J, Cybulski C, Oszurek O, et al. Serum selenium level and cancer risk: a nested case-control study. *Hereditary cancer in clinical practice*. 2019;17(1):1-7. doi:10.1186/s13053-019-0131-7 PMID:31890061
 - Kristal AR, Darke AK, Morris JS, Tangen CM, Goodman PJ, Thompson IM, et al. Baseline selenium status and effects of selenium and vitamin E supplementation on prostate cancer risk. *Journal of the National Cancer Institute*. 2014;106(3):djt456. doi:10.1093/jnci/djt456
 - Semnani Sh, Roshandel Gh, Keshkar AA, Zendeabad A, Rahimzadeh H, Besharat S, et al. Relationship between Soil Selenium level and esophageal cancer: An ecological study in Golestan province of Iran. *Journal of Gorgan University of Medical Sciences*. 2010;12(3):51-6. doi:10.1016/j.jtemb.2010.03.002 PMID:20569930
 - Jaworska K, Gupta S, Durda K, Muszyńska M, Sukiennicki G, Jaworowska E, et al. A low selenium level is associated with lung and laryngeal cancers. *PLoS one*. 2013;8(3):e59051. doi:10.1371/journal.pone.0059051 PMID:23516596